

AUTHOR INDEX

- Andersson, F.O. 149
Andrews, M.A. 141
- Benefice-Malouet, S. 293
Bischoff, M. 17
Blanc-Muesser, M. 225
Bock, K. 213
Brade, H. 213
Byrd, R.A. 195
- Christian, R. 213
Coe, P.L. 293
- Driguez, H. 225
- Egan, W. 195
- Furui, H. C1
- Glaudemans, C.P.J. 103, 271
Goldstein, I.J. 337, 347
Gómez-Sánchez, A. 307
Gould, G.L. 141
Grand-Maitre, C. 149
Grasdalen, H. 233
Greiner, J. 323
- Hadjoudis, E. 1
Hasegawa, A. C1
Hashimoto, H. C5
Hatfield, R.D. 183
Helm, R.F. 183
Hermosín, I. 307
Hisamichi, K. 369
Holst, O. 213
- Iiyama, K. 131
Ishizu, A. 131
- Jaekel, F. 167
- Kajihara, Y. C5
Kaku, H. 337, 347
- Kamerling, J.P. 155
Kasuya, N. 131
Kiso, M. C1
Knutsen, S.H. 233
Kobayashi, H. 369
Kobayashi, Y. 363
Kodama, H. C5
Koga, K. 363
Köll, P. 17
Kong, F. 75
Konno, S. 369
Kopf, J. 17
Kosma, P. 213
Kováč, P. 103
- Lankin, D.C. 245
Lavaitte, S. 225
Lis, T. 33
Lu, D. 75
Lukacs, G. 161, 167
- Mavridis, I.M. 1
Maya, I. 307
Mazeau, K. 225
Milius, A. 323
Morf, M. 17
- Nagumo, T. 355
Nishi, K. 117, 173
Nishino, T. 355
Nugent, S.T. 245
- Ogawa, K. 41, 57
Ogawa, S. 117, 173
Olesker, A. 161, 167
- Pavliak, V. 103
Peumans, W.J. 347
- Ralph, J. 183
Rao, S.N. 245
Rémy, S. 167
Riess, J.G. 323
Roy, R. 149

Sarda, P. 161
Sarko, A. 41, 57
Shibata, N. 369
Somogyi, L. 89
Sutoris, H. 271
Suzuki, S. 369

Taneichi, N. 363
Taravel, F.R. 225
Thomsen, J.U. 213
Tonegawa, T. 173
Troppe, F.D. 149
Tsuchiya, T. 363
Tvaroska, I. 225

Van Damme, E.J.M. 347
Van der Ven, J.G.M. 155

Van Seeventer, P. 155
Van Steijn, A.M.P. 155
Vliegthart, J.F.G. 155

Walker, R.T. 293
Widmalm, G. 195
Wu, X. 75

Yokoyama, J. 173
Yui, T. 41, 57

Zhang, P. 75
Ziegler, T. 271
Zimmer, B. 17

SUBJECT INDEX

- 2-*O*-(2-Acetamido-2-deoxy- β -D-glucopyranosyl)-5a-carba- α -D-mannopyranose and 5a-carba-2-*O*- and 3,6-di-*O*-(α -D-mannopyranosyl)- α -D-mannopyranoses, synthesis of, 117
- 2-*C*-Acetamidomethyl-2-deoxy- α -D-glucopyranoside and its *manno* isomer, synthesis of methyl, 161
- 3-(1-Acetamidopropyl)pyrazolo[3,4-*b*]quinoxaline (flavazole) derivatives, preparation of, 89
- Allofuranose derivatives, NMR spectroscopy and conformational analysis of substituted 1,2:5,6-di-*O*-isopropylidene- α -D-, 245
- Amadori compounds, influence of malondialdehyde on the Maillard degradation of, 307
- Anionic surfactant for biomedical use, synthesis of D-glucose 3- and 6-[2-(perfluoroalkyl)ethyl phosphates] as a new type of, 323
- Anticoagulant and antithrombin activities of oversulfated fucans, 355
- Binding studies on immunodeterminants: synthesis of β -(1 \rightarrow 6)-linked oligosaccharide methyl glycosides having 1–4 internal D-galactopyranosyl residues flanked by gentiobiose residues, 271
- Building blocks for carba-oligosaccharides related to cell-surface glycans, preparation of, 173
- 5a-Carba-2-*O*- and 3,6-di-*O*-(α -D-mannopyranosyl)- α -D-mannopyranoses, synthesis of 2-*O*-(2-acetamido-2-deoxy- β -D-glucopyranosyl)-5a-carba- α -D-mannopyranose and, 117
- Carba-oligosaccharides related to cell-surface glycans, preparation of building blocks for, 173
- Carbon–proton coupling constants for the C–S–C–H pathway in 1-thioglycosides, Karplus-type equation for vicinal, 225
- Chiral polysubstituted *cis*-fused decalin containing an angular C-methyl group, synthesis by an intramolecular Diels–Alder reaction, 167
- Chlamydia* lipopolysaccharides, an NMR spectroscopic investigation of Kdo-containing oligosaccharides related to the genus-specific epitope of, 213
- Conformational analysis of substituted 1,2:5,6-di-*O*-isopropylidene- α -D-allofuranose derivatives, NMR spectroscopy and, 245
- Conformational study of α -L-Rhap-(1 \rightarrow 2)- α -L-Rhap-(1 \rightarrow OMe) by NMR nuclear Overhauser effect spectroscopy (NOESY) and molecular dynamics calculations, 195
- p*-Coumaroylated and feruloylated methyl glycosides, synthesis of, 183
- Crystal and molecular structures of the hexitol hexa-acetates, 17
- Crystal structure of α -D-glucopyranosyl potassium hydrogenphosphate, 33
- Crystal structure of konjac glucomannan in the mannan II polymorphic form, molecular and, 41
- Crystal structure of the inclusion complex of cyclomaltoheptaose (β -cyclodextrin) with 4-*tert*-butyltoluene, 1
- Crystal structure of the regenerated form of (1 \rightarrow 3)- α -D-mannan, molecular and, 57
- Cyclomaltoheptaose (β -cyclodextrin), crystal structure of the inclusion complex of 4-*tert*-butyltoluene with, 1
- Decalin containing an angular C-methyl group, synthesis of a chiral polysubstituted *cis*-fused, by an intramolecular Diels–Alder reaction, 167
- Deoxyfluorocellulose acetate, synthesis and characterization of highly substituted, 131
- 1-Deoxynojirimycin-containing glycans related to the Lewis X and sialyl-Lewis X epitopes recognized by LEC-CAMs, synthesis of, c1
- Diels–Alder reaction, synthesis of a chiral polysubstituted *cis*-fused decalin containing an angular C-methyl group by an intramolecular, 167

- 2,6-Dioxabicyclo[3.1.1]heptanes, substituted: synthesis and conformational analysis of 1,3-anhydro-2,4-di-*O*-benzyl-6-deoxy- β -D-glucopyranose, 75
- Disaccharides in aqueous sodium hydroxide, epimerization of reducing terminal groups of (1 \rightarrow 2)-linked D-gluco- and D-manno-, 369
- Epimerization of reducing terminal groups of (1 \rightarrow 2)-linked D-gluco- and D-manno-disaccharides in aqueous sodium hydroxide, 369
- Feruloylated and *p*-coumaroylated methyl glycosides, synthesis of, 183
- Fucans, anticoagulant and antithrombin activities of oversulfated, 355
- D-Galactopyranosyl residues flanked by gentiobiose residues, synthesis of β -(1 \rightarrow 6)-linked oligosaccharide methyl glycosides having 1–4 internal, 271
- Garlic (*Allium sativum*) bulbs, new mannose-specific lectin from, 347
- β -D-Glucopyranose, 1,3-anhydro-2,4-di-*O*-benzyl-6-deoxy, synthesis, and conformational analysis by ^1H -NMR spectroscopy and molecular mechanics calculation, 75
- α -D-Glucopyranosyl potassium hydrogenphosphate, the crystal structure of, 33
- D-Glucose 3- and 6-[2-(perfluoroalkyl)ethyl phosphates], synthesis of a new type of anionic surfactant for biomedical use, 323
- β -D-Glycobiosides, 1-thio-, 4-nitrophenyl, phase transfer catalyzed synthesis of, 149
- Glycosides, methyl, synthesis of feruloylated and *p*-coumaroylated, 183
- Hexitol hexa-acetates, the crystal and molecular structures of the, 17
- Inclusion complex of cyclomaltoheptaose (β -cyclodextrin) with 4-*tert*-butyltoluene, crystal structure of the, 1
- Isopropylidene and benzylidene derivatives of sugars: transacetalation, a convenient non-aqueous method for effecting the deprotection of, 141
- Karplus-type equation for vicinal carbon–proton coupling constants for the C–S–C–H pathway in 1-thioglycosides, 225
- Kdo-containing oligosaccharides related to the genus-specific epitope of *Chlamydia* lipopolysaccharides, an NMR spectroscopic investigation of, 213
- Konjac glucomannan in the mannan II polymorphic form, molecular and crystal structure of, 41
- Lectins from bulbs, mannose-specific, interaction with linear manno-oligosaccharides, 337
- Lewis X and sialyl-Lewis X epitopes, synthesis of related 1-deoxynojirimycin-containing glycans, C1
- Linked D-gluco- and D-manno-disaccharides in aqueous sodium hydroxide, epimerization of reducing terminal groups of (1 \rightarrow 2)-, 369
- Maillard degradation of Amadori compounds, influence of malondialdehyde on the, 307
- Malondialdehyde, influence on the Maillard degradation of Amadori compounds, 307
- (1 \rightarrow 3)- α -D-Mannan, molecular and crystal structure of the regenerated form of, 57
- Mannose-specific bulb lectins, interaction with linear manno-oligosaccharides and comparison with mannose/glucose-binding lectins, 337
- Mannose-specific lectins from garlic and ramsons bulbs, 347
- Methyl 3-*O*-(2-acetamido-2-deoxy-6-thio- β -D-glucopyranosyl)- β -D-galactopyranoside: a slow reacting acceptor-analogue which inhibits glycosylation by UDP-D-galactose-*N*-acetyl-D-glucosamine-(1 \rightarrow 4)-D-galactosyl-transferase, c5
- Methyl 2-*C*-acetamidomethyl-2-deoxy- α -D-glucopyranoside and its *manno* isomer, synthesis of, 161
- Methyl 3-deoxy-3-*C*-formyl- α -D-*arabino*-pentofuranoside, synthesis of derivatives of, 293
- Molecular and crystal structure of konjac glucomannan in the mannan II polymorphic form, 41
- Molecular and crystal structures of the hexitol hexa-acetates, 17
- Molecular and crystal structure of the regenerated form of (1 \rightarrow 3)- α -D-mannan, 57
- Neocarrabiose oligosaccharides having different length and sulphate substitution, use as model for ^1H -NMR spectroscopy, 233
- 4-Nitrophenyl 1-thio- β -D-glycobiosides, phase transfer synthesis of, 149
- NMR NOESY spectroscopy and molecular dynamics calculations, conformational study of α -L-Rha *p*-(1 \rightarrow 2)- α -L-Rha *p*-(1 \rightarrow OMe) by, 195
- NMR spectroscopic investigation of Kdo-containing oligosaccharides related to the genus-specific epitope of *Chlamydia* lipopolysaccharides, 213

- NMR spectroscopy and conformational analysis of substituted 1,2:5,6-di-*O*-isopropylidene- α -D-allofuranose derivatives, 245
- NMR spectroscopy, the use of neocarrabiose oligosaccharides having different length and sulphate substitution as model compounds for ^1H -, 233
- Oligosaccharide fragments of *Shigella dysenteriae* type 1 O-antigen, stereoselective syntheses of, 103
- Oligosaccharide methyl glycosides having 1–4 internal D-galactopyranosyl residues flanked by gentiobiose residues, synthesis of β -(1 \rightarrow 6)-linked, for immunodeterminant binding studies, 271
- Oligosaccharides having different length and sulphate substitution as model compounds for ^1H -NMR spectroscopy, the use of neocarrabiose, 233
- Oversulfated fucans, anticoagulant and antithrombin activities of, 355
- Pentasaccharide fragment of the capsular polysaccharide of *Streptococcus pneumoniae* type 18C, synthesis of a repeating, 155
- Pentofuranoside, synthesis of derivatives of methyl 3-deoxy-3-*C*-formyl- α -D-*arabino*-, 293
- Phenylhydrazone derivatives, addition and cyclisation reactions of sugar, via a 1,4-elimination process, 89
- Polysaccharide of *Streptococcus pneumoniae* type 18C, synthesis of a repeating pentasaccharide fragment of the capsular, 155
- Potassium hydrogenfluoride, reaction of 2-deoxy-6-*O*-[2,3-dideoxy-4,6-*O*-isopropylidene-2,3-(*N*-tosylepimino)- α -D-mannopyranosyl]-4,5-*O*-isopropylidene-1,3-di-*N*-tosylstreptamine with, 363
- Preparation of building blocks for carba-oligosaccharides related to cell-surface glycans, 173
- Quinoxalinone derivatives, preparation of 3-(2-acetamido-1-phenylhydrazonobutyl)-2(1*H*)-, 89
- Ramsons (*Allium ursinum*) bulbs, new mannose-specific lectin from, 347
- Reaction of 2-deoxy-6-*O*-[2,3-dideoxy-4,6-*O*-isopropylidene-2,3-(*N*-tosylepimino)- α -D-mannopyranosyl]-4,5-*O*-isopropylidene-2,3-di-*N*-tosylstreptamine with potassium hydrogenfluoride, 363
- α -L-Rhap-(1 \rightarrow 2)- α -L-Rhap-(1 \rightarrow OMe), conformational study by NMR NOESY spectroscopy and molecular dynamics calculations, 195
- Shigella dysenteriae* type 1 O-antigen, stereoselective syntheses of a di-, tri-, and tetra-saccharide fragment of, 103
- Sialyl-Lewis X and Lewis X epitopes, synthesis of related 1-deoxynojirimycin-containing glycans, C1
- Streptococcus pneumoniae* type 18C, synthesis of a repeating pentasaccharide fragment of the capsular polysaccharide of, 155
- Surfactant for biomedical use, synthesis of D-glucose 3- and 6-[2-(perfluoroalkyl)ethyl phosphates] as a new type of anionic, 323
- Synthesis and characterization of highly substituted deoxyfluorocellulose acetate, 131
- Synthesis of 2-*O*-(2-acetamido-2-deoxy- β -D-glucopyranosyl)-5a-carba- α -D-mannopyranose and 5a-carba-2-*O*- and 3,6-di-*O*-(α -D-mannopyranosyl)- α -D-mannopyranoses, 117
- Synthesis of derivatives of methyl 3-deoxy-3-*C*-formyl- α -D-*arabino*-pentofuranoside, 293
- Synthesis of methyl 2-*C*-acetamidomethyl-2-deoxy- α -D-glucopyranoside and its *manno* isomer, 161
- 1-Thio- β -D-glycobiosides, 4-nitrophenyl, phase transfer catalyzed synthesis of, 149
- 1-Thioglycosides, Karplus-type equation for vicinal carbon-proton coupling constants for the C-S-C-H pathway in, 225
- Tosylepimino)- α -D-mannopyranosyl]-4,5-*O*-isopropylidene-1,3-di-*N*-tosylstreptamine with potassium hydrogenfluoride, reaction of 2-deoxy-6-*O*-[2,3-dideoxy-4,6-*O*-isopropylidene-2,3-(*N*-, 363
- Transacetalation: a convenient nonaqueous method for effecting the deprotection of isopropylidene and benzylidene derivatives of sugars, 141
- 3,4,6-Tri-*O*-acetyl-2-azido-2-deoxy- α -D-glucopyranosyl chloride, use as a glycosyl donor in stereoselective syntheses of fragments of *Shigella dysenteriae* type 1 O-antigen, 103

